

Recombinant Human Activin receptor
type-2B(ACVR2B),partial



Catalog No: #AP73748

Orders: order@abscitech.com

Package Size: #AP73748-1 10ug #AP73748-2 50ug #AP73748-3 100ug #AP73748-4 200ug #AP73748-5 500ug #AP73748-6 1mg

Support: tech@abscitech.com


Description

Product Name	Recombinant Human Activin receptor type-2B(ACVR2B),partial
Brief Description	Recombinant Protein
Host Species	Yeast
Target Name	ACVR2B
Other Names	Activin receptor type IIB Short name: ACTR-IIB
Accession No.	Uniprot ID: Q13705
Target Species	Hu
SDS-PAGE MW	15.66kDa
Target Length	Extracellular Domain,19-137aa
Tag Info	N-terminal 6xHis-tagged
Target Sequence	SGRGEAETRECIYYNANWELERTNQSGLERCEGEQDKRLHCYASWRNSSGTIELVKKG CWLDDFN CYDRQE CVATEENPQVYFCCCEGNFCNERFTHLPEAGGPEVTYEPPTAPTLLT
Formulation	Tris-based buffer,50% glycerol
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.

Application Details

Greater than 90% as determined by SDS-PAGE.

Images



Background

Transmembrane serine/threonine kinase activin type-2 receptor forming an activin receptor complex with activin type-1 serine/threonine kinase receptors (ACVR1, ACVR1B or ACVR1c). Transduces the activin signal from the cell surface to the cytoplasm and is thus regulating many physiological and pathological processes including neuronal differentiation and neuronal survival, hair follicle development and cycling, FSH production by the pituitary gland, wound healing, Extracellular domain matrix production, immunosuppression and carcinogenesis. Activin is also thought to have a paracrine or autocrine role in follicular development in the ovary. Within the receptor complex, the type-2 receptors act as a primary activin receptors (binds activin-A/INHBA, activin-B/INHBB as well as inhibin-A/INHA-INHBA). The type-1 receptors like ACVR1B act as downstream transducers of activin signals. Activin binds to type-2 receptor at the plasma membrane and activates its serine-threonine kinase. The activated receptor type-2 then phosphorylates and activates the type-1 receptor. Once activated, the type-1 receptor binds and phosphorylates the SMAD proteins SMAD2 and SMAD3, on serine residues of the C-terminal tail. Soon after their association with the activin receptor and subsequent phosphorylation, SMAD2 and SMAD3 are released into the cytoplasm where they interact with the common partner SMAD4. This SMAD complex translocates into the nucleus where it mediates activin-induced transcription. Inhibitory SMAD7, which is recruited to ACVR1B through FKBP1A, can prevent the association of SMAD2 and SMAD3 with the activin receptor complex, thereby blocking the activin signal. Activin signal transduction is also antagonized by the binding to the receptor of inhibin-B via the IGSF1 inhibin coreceptor.

References

"Expression of type II activin receptor genes during differentiation of human K562 cells and cDNA cloning of the human type IIB activin receptor."Hilden K., Tuuri T., Eramaa M., Ritvos O.Blood 83:2163-2170(1994)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.