Tau(Phospho-Thr181) Antibody

Catalog No: #AB11107

Package Size: #AB11107-1 50ul #AB11107-2 100ul #AB11107-4 25ul




Orders: order@abscitech.com Support: tech@abscitech.com

Description Tau(Phospho-Thr181) Antibody Product Name Host Species Rabbit Clonality Polyclonal Purification Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy using non-phosphopeptide. WB IHC Applications Species Reactivity Hu Ms Rt Specificity The antibody detects endogenous level of Tau only when phosphorylated at threonine 181. Peptide-KLH Immunogen Type Peptide sequence around phosphorylation site of threonine 181 (P-K-T(p)-P-P) derived from Human Tau. Immunogen Description Target Name Tau Modification Phospho-Thr181 Other Names MAPT; MTAPT; MTBT1; Neurofibrillary tangle protein; PHF-tau Accession No. Swiss-Prot: P10636NCBI Protein: NP _001116538.1 Concentration 1.0mg/ml Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Storage Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 48 62 78 kd Western blotting: 1:500~1:1000 Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from mouse brain tissue using Tau(Phospho-Thr181) Antibody #AB11107 and the same antibody preincubated with blocking peptide.



Immunohistochemical analysis of paraffin-embedded rat hippocampal region tissue from a model with Alzheimer

Background

Ρ

Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Puig B, et al.(2005) Acta Neuropathol (Berl). 110(3):261-268.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.