Mouse anti-Human CD8, APC Conjugated mAb

Catalog No: #AB28029

Package Size: #AB28029-1 50 Tests



Orders: order@abscitech.com Support: tech@abscitech.com

Description


Product Name	Mouse anti-Human CD8, APC Conjugated mAb
Host Species	Mouse
Clonality	Monoclonal
Clone No.	COV8
Isotype	lgG1
Applications	FC
Species Reactivity	Hu
Conjugates	APC
Target Name	CD8
Formulation	Phosphate-buffered solution, pH7.4, containing 0.09% sodium azide and 0.2% (w/v) BSA
Storage	Store at 4°C. DO NOT FREEZE. LIGHT SENSITIVE MATERIAL.

Application Details

Vol.per.Test: 10 μl/Test

Notice: This reagent has been pre-diluted for use at recommended volume per test in flow cytometry analysis. Typically add 10ul of this reagent to 100μl of experimental sample with 1 X 106 cells per test. Please refer to the detailed protocol when you perform a test.

Images



Human peripheral blood lymphocytes analyzed with APC CD8 mAb

Background

COV8 reacts with the 32 kDa a chain of CD8. This molecule is expressed on the T suppressor/cytotoxic cell population (which comprises about 1/3 of the peripheral blood T lymphocytes total population) and with most of thymocytes, as well as a subset of NK cells. CD8 expresses as either a heterodimer with the CD8b chain (CD8ab) or as a homodimer (CD8aa or CD8bb). CD8 acts as a co-receptor with MHC Class I restricted TCRs in antigen recognition. CD8 function is important for positive selection of MHC Class I restricted CD8+ T cells during T cell development.

- * Ratio of CD4 and CD8 is important for monitor of HIV infection.
- 1. Engleman EG et al. (1981) J Exp Med 153: 193-198.
- 2. Haynes, B et al. (1986) Leukocyte Typing II: Human T lymphocytes. New York: Sprimger-Verlag; 3-30.

- 3. Parnes, JR. et al. (1989). Adv. Immunol. 44,256-311.
- 4. Bikah G. et al. (1996). Science 274:1906.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.